

Process $3 \rightarrow 3$ and crossing symmetry violation

E. Bartoš^{1,2}, S. R. Gevorkyan^{1,3}, E. A. Kuraev¹

¹ *Joint Institute of Nuclear Research, 141980 Dubna, Russia*

² *Dept. of Theor. Physics, Comenius Univ., 84248 Bratislava, Slovakia*

³ *Yerevan Physics Institute, 375036 Yerevan, Armenia*

Abstract

Using the Sudakov technique we sum the perturbation series for the process $3 \rightarrow 3$ and obtain the compact analytical expression for the amplitude of this process, which takes into account all possible Coulomb interactions between colliding particles. Compare it with the amplitude of the lepton pair production in heavy ion collision i.e. in the process $2 \rightarrow 4$, we show that crossing symmetry between this processes holds only if one neglects the interaction of produced pair with ions (i.e. in the approximation $Z_{1,2}\alpha \ll 1$).

1 Introduction

During the past decade the growing interest to the process of lepton pair production in the strong Coulomb fields appeared. This is connected mainly with beginning of operation the relativistic heavy ion collider RHIC (Lorentz factor $\gamma = \frac{E}{m} = 108$) and the new collider LHC ($\gamma = 3000$) which will operate in the nearest future. At such energies the lepton pairs yield becomes huge (remind that the total cross section of the process $A + B \rightarrow A + B + e^+ + e^-$ grows with energy as the third power of the logarithm [1, 2]), so that a detail analysis of this process, with taking into account the Coulomb corrections (CC) is required. Such work has been done during last years and a lot of

papers are devoted to this subject. Nevertheless the problem turn out to be more complex than it seems from the first glance. For a review of the works in this direction see for example [3] and references therein. We want only to notice the exciting result obtained in the series of papers [4]: the Coulomb corrections to the process $A + B \rightarrow A + B + e^+ + e^-$ enters the amplitude of this process in such a way that its cross section is determined solely by lowest (Born) term. At present we understand that this result is the incorrect application of crossing symmetry property which, as it is known long ago (see e.g. [5]), is valid only on the Born level. As an obvious example of the crossing symmetry violation we want to cite the process of lepton pair photoproduction on the nuclei and its counterpart the bremsstrahlung in lepton–nucleus scattering. Amplitudes of both processes are determined by Coulomb phase which is infrared stable in the case of pair photoproduction while it is infrared divergent in the case of bremsstrahlung and this difference can't be adjust by trivial crossing change of variables. Nevertheless taking into account the importance of the problem and permanent interest to it from scientific society we calculate the full amplitude for the process $3 \rightarrow 3$ accounting all possible photon exchanges among the colliding relativistic particles. From its comparison with the amplitude of the process $2 \rightarrow 4$ one can see that the crossing symmetry property becomes invalid whereas one takes into account the final state interaction.

2 The Born amplitude

Let us construct the amplitude of the process $3 \rightarrow 3$ represented in Fig. 1 (a, b)

$$A_1(p_1) + A_2(p_2) + C(p_3) \rightarrow A_1(p'_1) + A_2(p'_2) + C(p'_3). \quad (1)$$

We consider the kinematics when all the energy invariants which determined the process (1) are large, compared with the masses of involved particles and the transferred momenta

$$\begin{aligned} s &= (p_1 + p_2)^2, & s_1 &= (p_1 + p_3)^2, & s_2 &= (p_3 + p_2)^2, \\ q_1^2 &= (p_1 - p'_1)^2, & q_2^2 &= (p_2 - p'_2)^2, & q_3^2 &= (p_3 - p'_3)^2, \\ p_1^2 &= p'^2_1 = m_1^2, & p_2^2 &= p'^2_2 = m_2^2, & p_3^2 &= p'^2_3 = m^2, \\ s &\gg s_1 \sim s_2 \gg -q_1^2 \sim -q_2^2 \sim -q_3^2. \end{aligned} \quad (2)$$

For the Born amplitude of the process (1) one can write

$$M_{(1)}^{(1)} = -i(4\pi\alpha)^2 Z_1 Z_2 \bar{u}(p'_1) \gamma_\mu u(p_1) \bar{u}(p'_2) \gamma_\nu u(p_2) \frac{\bar{u}(p'_3) O_{\rho\sigma} u(p_3) g^{\mu\sigma} g^{\nu\rho}}{q_1^2 q_2^2}, \quad (3)$$




Fig. 1: Feynman diagrams for Born amplitudes of the process $A_1 + A_2 + C \rightarrow A_1 + A_2 + C$ (a, b) and the process $A_1 + A_2 \rightarrow A_1 + A_2 + C + \bar{C}$ (c, d).

where $Z_{1,2}$ are the charge numbers of the colliding nuclei. We use Sudakov's parameterization for all four-momenta entering the problem (for details see [6])

$$\begin{aligned} q_1 &= \alpha_1 \tilde{p}_2 + \beta_1 \tilde{p}_1 + q_{1\perp}, & q_2 &= \alpha_2 \tilde{p}_2 + \beta_2 \tilde{p}_1 + q_{2\perp}, \\ p'_1 &= \alpha'_1 \tilde{p}_2 + \beta'_1 \tilde{p}_1 + p'_{1\perp}, & p'_2 &= \alpha'_2 \tilde{p}_1 + \beta'_2 \tilde{p}_1 + p'_{2\perp}, \\ p_3 &= \alpha_3 \tilde{p}_1 + \beta_3 \tilde{p}_1 + p_{3\perp}, & p'_3 &= \alpha'_3 \tilde{p}_1 + \beta'_3 \tilde{p}_1 + p'_{3\perp}, \end{aligned} \quad (4)$$

and the Gribov's decomposition of the metric tensor in the longitudinal and transverse parts

$$g_{\mu\nu} = g_{\perp\mu\nu} + \frac{2}{s} (\tilde{p}_{1\mu} \tilde{p}_{2\nu} + \tilde{p}_{1\nu} \tilde{p}_{2\mu}),$$

with light-like 4-vectors $\tilde{p}_{1,2}$. For the kinematics of the process we have

$$\begin{aligned} s &= 2\tilde{p}_1 \tilde{p}_2, & \beta_1 + \beta_3 &= \beta'_3, & \alpha_2 + \alpha_3 &= \alpha'_3, \\ g_{\mu\sigma} &= \frac{2}{s} \tilde{p}_{1\sigma} \tilde{p}_{2\mu}, & g_{\nu\rho} &= \frac{2}{s} \tilde{p}_{1\nu} \tilde{p}_{2\rho}, \\ q_1^2 &= q_{1\perp}^2 = -\mathbf{q}_1^2, & q_2^2 &= q_{2\perp}^2 = -\mathbf{q}_2^2, \end{aligned} \quad (5)$$

where we use two-dimensional vectors in the plane transverse to the z-axes, which we choose along 3-vector $\vec{p}_1 = -\vec{p}_2$ in the center of mass frame of initial particles A_1, A_2 . Using the gauge invariance conditions

$$\begin{aligned} q_{1\rho} \bar{u}(p'_3) O_{\rho\sigma} u(p_3) &\approx (\beta_1 \tilde{p}_1 + q_{1\perp})_\rho \bar{u}(p'_3) O_{\rho\sigma} u(p_3) = 0, \\ q_{2\sigma} \bar{u}(p'_3) O_{\rho\sigma} u(p_3) &\approx (\alpha_2 \tilde{p}_2 + q_{2\perp})_\sigma \bar{u}(p'_3) O_{\rho\sigma} u(p_3) = 0, \end{aligned} \quad (6)$$

we get the Born amplitude in the form

$$M_{(1)}^{(1)}(q_1, q_2) = -4isN_1N_2(4\pi\alpha Z_1)(4\pi\alpha Z_2)B(q_1, q_2), \quad (7)$$




Fig. 2: Feynman diagrams for the process $A_1 + A_2 + C \rightarrow A_1 + A_2 + C$ with three photon exchange.

with

$$B(q_1, q_2) = \frac{\bar{u}(p'_3) O_{\rho\sigma} u(p_3) q_{1\perp}^\rho q_{2\perp}^\sigma}{s\alpha_2\beta_1 \mathbf{q}_1^2 \mathbf{q}_2^2}, \quad (8)$$

$$N_1 = \frac{1}{s} \bar{u}(p'_1) \hat{p}_2 u(p_1), \quad N_2 = \frac{1}{s} \bar{u}(p'_2) \hat{p}_1 u(p_2),$$

$$s\alpha_2\beta_1 = -q_3^2 - (\mathbf{q}_1 - \mathbf{q}_2)^2 \sim m^2.$$

The values of N_i for every polarization state of initial particles (or for spinless particles) are unity and

$$\bar{u}(p'_3) O_{\rho\sigma} u(p_3) q_{1\perp}^\rho q_{2\perp}^\sigma = \bar{u}(p'_3) \left[\hat{q}_{2\perp} \frac{\hat{p}_3 + \hat{q}_1 + m}{(p_3 + q_1)^2 - m^2} \hat{q}_{1\perp} + \hat{q}_{1\perp} \frac{\hat{p}_3 + \hat{q}_2 + m}{(p_3 + q_2)^2 - m^2} \hat{q}_{2\perp} \right] u(p_3). \quad (9)$$

3 The Coulomb corrections to the process $3 \rightarrow 3$

Let us consider the set of six Feynman diagrams (FD) with one virtual photon connected the p_3 line with the particle A_1 and the two ones connected p_3 line with the particle A_2 (see Fig. 2). The loop momentum integration in the




Fig. 3: Feynman diagram for the amplitude $M_{(2)}^{(1)}$.

relevant matrix element can be performed accounting that

$$d^4k = (2\pi i)^2 \frac{1}{2s} \frac{d(s\alpha_k)}{2\pi i} \frac{d(s\beta_k)}{2\pi i} d^2k_\perp, \quad k = \alpha_k \tilde{p}_2 + \beta_k \tilde{p}_1 + k_\perp. \quad (10)$$

It can be shown that only 4 FD amplitudes works (Fig. 2 (a–d)). Really, when one performs the α_k integration in Fig. 2 (e, f), one has a situation when both poles in α_k complex plane are situated in the same half-plane, so their contribution to the amplitude is zero (suppressed by factor $|q_3^2/s| \sim |s_1/s|$).

The physical reason of this suppression is the same as in the case of bremsstrahlung suppression for fast charged particle moving through the media known as the Landau–Pomeranchuk effect. Really, this effect can be explained starting from the fact of power suppression of radiation between two scattering centers in the case when the distance between these centers is less than the coherent length.

It is convenient to introduce 8 FD (including remaining four ones and additional four FD with interchanged photons absorbed by nucleus A_2 line). To avoid the double counting we multiply the relevant matrix element by statistical factor $1/2!$. This trick permits to perform the integration over Sudakov variables α_k, β_k with the result

$$\int \frac{d(s\alpha_k)}{2\pi i} \left(\frac{1}{s\alpha_k + i0} + \frac{1}{-s\alpha_k + i0} \right) = \int \frac{d(s\beta_k)}{2\pi i} \left(\frac{1}{s\beta_k + i0} + \frac{1}{-s\beta_k + i0} \right) = 1. \quad (11)$$

Further integration over transverse momentum is straightforward

$$\int \frac{d^2\mathbf{k}}{\pi} \frac{1}{(\mathbf{k} + \lambda)((\mathbf{q}_2 - \mathbf{k})^2 + \lambda^2)} = \frac{2}{\mathbf{q}_2^2} \ln \frac{\mathbf{q}_2^2}{\lambda^2}. \quad (12)$$




Fig. 4: Feynman diagram for the amplitude $M_{(\infty)}^{(\infty)}(q_1, q_2)$.

For the amplitude $M_{(2)}^{(1)}$ (see Fig. 3) and the similar amplitude $M_{(1)}^{(2)}$ we obtained

$$M_{(2)}^{(1)} + M_{(1)}^{(2)} = M_{(1)}^{(1)}(q_1, q_2) \frac{1}{2!} \left[2iZ_1\alpha \ln \frac{\mathbf{q}_1^2}{\lambda^2} + 2iZ_2\alpha \ln \frac{\mathbf{q}_2^2}{\lambda^2} \right]. \quad (13)$$

The amplitude for arbitrary amount of interchanged photons (see Fig. 4) is constructed in the similar way

$$M_{(\infty)}^{(\infty)}(q_1, q_2) = M_{(1)}^{(1)}(q_1, q_2) e^{i(\varphi_1(\mathbf{q}_1) + \varphi_2(\mathbf{q}_2))} \quad (14)$$

with the Coulomb phases

$$\varphi_1(\mathbf{q}_1) = Z_1\alpha \ln \frac{\mathbf{q}_1^2}{\lambda^2}, \quad \varphi_2(\mathbf{q}_2) = Z_2\alpha \ln \frac{\mathbf{q}_2^2}{\lambda^2}.$$

Consider now the case with one additional exchanged photon between two nuclei A_1, A_2 . The relevant matrix element $M_{(1B)}$ reads

$$M_{(1B)} = i\alpha Z_1 Z_2 \int \frac{d^2\mathbf{k}}{\pi(\mathbf{k}^2 + \lambda^2)} M_{(1)}^{(1)}(q_1 + k, q_2 + k). \quad (15)$$

Two-dimensional integral in (15) is infrared divergent. To regularize it we introduce the photon mass parameter λ .

In the same approach we get for the matrix element with the n exchanged (between nuclei) photons (see Fig. 5 (a))

$$M_{(nB)} = \frac{(i\alpha Z_1 Z_2)^n}{n!} \prod_{i=1}^n \int \frac{d^2\mathbf{k}_i}{\pi(\mathbf{k}_i^2 + \lambda^2)} M_{(1)}^{(1)}(q_1 + \sum_{i=1}^n k_i, q_2 + \sum_{i=1}^n k_i). \quad (16)$$




Fig. 5: Feynman diagrams for the n photon exchange between nuclei A_1 and A_2 compared with the Born diagram for the processes $3 \rightarrow 3$ (a) and $2 \rightarrow 4$ (b) (blob in (b) correspond to diagrams in Fig. 1 (c, d)) .

It is convenient to write down this expression in impact parameter representation. For this aim we use the following identity

$$\int d^2 \mathbf{k}_{n+1} \delta^{(2)}(k_{n+1} - q_1 - \sum_{i=1}^n k_i) = \frac{1}{(2\pi)^2} \int d^2 \mathbf{k}_{n+1} d^2 \boldsymbol{\rho} e^{i(\mathbf{k}_{n+1} - \mathbf{q}_1 - \sum \mathbf{k}_i) \cdot \boldsymbol{\rho}} = 1. \quad (17)$$

Thus the matrix element with arbitrary number of exchanged photons can be cast

$$M(3 \rightarrow 3) = \sum_{n=1}^{\infty} M_{(nB)} = \frac{1}{4} \int \frac{d^2 \boldsymbol{\rho}}{\pi} e^{-i\mathbf{q}_1 \cdot \boldsymbol{\rho}} e^{i\alpha Z_1 Z_2 \psi(\boldsymbol{\rho})} \tilde{M}_{(1)}^{(1)}(\boldsymbol{\rho}, q_1, q_2) \quad (18)$$

with

$$\psi(\boldsymbol{\rho}) = \int \frac{d^2 \mathbf{k}}{\pi} \frac{e^{-i\mathbf{k} \cdot \boldsymbol{\rho}}}{\mathbf{k}^2 + \lambda^2} = 2K_0(\rho\lambda) = -2(\ln \rho m + \ln \frac{\lambda}{m} - \ln 2 + C), \quad (19)$$

where $C = 0.577$ and

$$\tilde{M}_{(1)}^{(1)}(\boldsymbol{\rho}, q_1, q_2) = \int \frac{d^2 \mathbf{k}}{\pi} e^{-i\mathbf{k} \cdot \boldsymbol{\rho}} M_{(1)}^{(1)}(k, k + q_2 - q_1). \quad (20)$$

This result confirms the general ansatz given above (see (14)) that all the dependence on "photon mass" λ can be represented as a phase factor. As can be seen the whole amplitude (18) cannot be cast solely as a Born amplitude

multiplied by the phase factor. The corresponding contributions to the total cross section (except the Born term) will be enhanced only by the first power of logarithm in energy.

Finally, taking into account all photon exchanges between particle C and nuclei A_1, A_2 we obtain the general answer by the simple replacement in the expression (20)

$$M_{(1)}^{(1)} \rightarrow M_{(\infty)}^{(\infty)} = M_{(1)}^{(1)}(k, k + q_2 - q_1) e^{i\varphi_1(\mathbf{k}) + i\varphi_2(\mathbf{k} + \mathbf{q}_2 - \mathbf{q}_1)} \quad (21)$$

with φ_1, φ_2 given in (14).

4 The Coulomb corrections to the process $2 \rightarrow 4$

As was mentioned above, our goal is to investigate the crossing symmetry property between the amplitudes of the process (1) and the relevant process in Fig. 1 (c, d)

$$A_1(p_1) + A_2(p_2) \rightarrow A_1(p'_1) + A_2(p'_2) + C(p_3) + \bar{C}(p_4), \quad (22)$$

with the following kinematics

$$\begin{aligned} s &= (p_1 + p_2)^2, \quad s_p = (q_+ + q_-)^2, \\ q_1^2 &= (p_1 - p'_1)^2, \quad q_2^2 = (p_2 - p'_2)^2, \\ p_1^2 &= p_1'^2 = m_1^2, \quad q_+^2 = q_-^2 = m^2, \quad q_+^2 = q_-^2 = m^2, \\ s &\gg -q_1^2 \sim -q_2^2 \sim s_{12}. \end{aligned} \quad (23)$$

Using the Sudakov technique the Born amplitude for the process (22) can be represented in the form

$$M_p = -is2^6\pi^2 Z_1 Z_2 N_1 N_2 B_p(q_1, q_2) \quad (24)$$

with

$$\begin{aligned} B_p(q_1, q_2) &= \frac{e_1^\alpha e_2^\beta \bar{u}(q_-) T_{\alpha\beta} v(q_+) |\mathbf{q}_1| |\mathbf{q}_2|}{\tilde{s} \mathbf{q}_1^2 \mathbf{q}_2^2}, \\ \tilde{s} &= s \alpha_2 \beta_1 = (q_+ + q_-)^2 + (\mathbf{q}_1 + \mathbf{q}_2)^2, \\ T_{\alpha\beta} &= \gamma_\beta \frac{\hat{q}_1 - \hat{q}_+ + m}{(q_1 - q_+)^2 - m^2} \gamma_\alpha + \gamma_\alpha \frac{\hat{q}_2 - \hat{q}_+ + m}{(q_2 - q_+)^2 - m^2} \gamma_\beta. \end{aligned}$$

(for details see [6]).

Generalization for the case of arbitrary number of exchanged photons between colliding nuclei is straightforward. The same approach as above leads to the following form of the generalized amplitude

$$M(2 \rightarrow 4) = \frac{1}{4} \int \frac{d^2 \boldsymbol{\rho}}{\pi} e^{-i \mathbf{q}_1 \boldsymbol{\rho} - i \alpha Z_1 Z_2 \psi(\boldsymbol{\rho})} \Phi_B(\boldsymbol{\rho}, q_2), \quad (25)$$

with

$$\Phi_B(\boldsymbol{\rho}, q_2) = \int \frac{d^2 \mathbf{k}}{\pi} e^{i \mathbf{k} \boldsymbol{\rho}} M_p(k, q_2 - k).$$

Comparing the expression (25) with the amplitude for the process $3 \rightarrow 3$ (18) one can see that the crossing symmetry property between the considered processes takes place in the case when one neglects the multiple exchanges of particle C with nuclei. Moreover, this statement is correct even when one takes into account the screening effects between nuclei A_1 and A_2 in both processes, which manifest itself by insertion of light-by-light scattering blocks into Feynman amplitudes. As was shown in [7] accounting of this effect can be provided by the universal factor

$$\exp \left\{ -\frac{\alpha^2 Z_1 Z_2}{2} L A(\rho) \right\}, \quad L = \ln(\gamma_1 \gamma_2), \quad (26)$$

with the complex quantity $A(\rho)$ connected with the Fourier transformation of light-by-light scattering amplitude and γ_1, γ_2 the Lorentz factors of colliding nuclei.

Nevertheless, crossing symmetry is broken in all orders of perturbation theory if one try to compare the full amplitude for the process $3 \rightarrow 3$ (expression (24) with the replacement (21)) and the relevant amplitude for the process $2 \rightarrow 4$ accounting the multiple interaction of produced particles [6].

Thus the crossing symmetry property takes place only for the colliding nuclei with charge numbers fulfilled the approximation $Z_{1,2} \alpha \ll 1$.

Acknowledgment

We are grateful to V. Serbo and N. Nikolaev for the useful discussions and comments. The work is supported by grants INTAS 97-30494, SR-2000.

References

- [1] L. D. Landau and E. M. Lifshits, Phys. Z. Sowjet. 6 (1934) 244.

- [2] G. Racah, *Nuovo Cimento* 14 (1937) 93.
- [3] R. N. Lee, A. I. Milstein, V. G. Serbo, hep-ph/0108014.
- [4] A. J. Baltz, L. McLerran, *Phys. Rev. C* 58, (1998) 1679.
- [5] E. A. Kuraev, L. N. Lipatov, M. I. Strikman, *Soviet ZhETP* 66 (1974) 838.
- [6] E. Bartoš, S. R. Gevorkyan, E. A. Kuraev and N. N. Nikolaev, hep-ph/0109281.
- [7] E. Bartoš, S. R. Gevorkyan, E. A. Kuraev and N. N. Nikolaev, hep-ph/0204327.
- [8] I Ginzburg, S. Panfil, V. Serbo, *Nucl. Phys. B* 284 (1987) 685.